arXiv:1101.1542v2 [astro-ph.HE] 21 Aug 2011

# Searching for curvature pion radiation from protons in strongly magnetized pulsars

**Douglas Fregolente**[1] **• Alberto Saa**[2]

**Abstract** By using rather conservative estimates based on the simplest polar cap model, we search the ATNF Pulsar Catalogue for strongly magnetized stars that could accelerate relativistic protons up to the curvature pion production threshold. The best candidate turns out to be the 16 ms pulsar J0537-6910, but the corresponding characteristic parameter $\chi = a/m_p$ is yet too small to give origin to observable signals. We show that, for pulsars with period $P \approx 1\,\mathrm{ms}$, a surface polar magnetic field $B \approx 10^{12}$G is required in order to induce detectable curvature pion radiation from accelerated protons in the magnetosphere. Some other emission processes are also considered.

**Keywords** pulsars, neutron stars, magnetic fields, high energy protons

## 1 Introduction

Processes related to the emission of pions by accelerated protons like, for instance, $p^+ \overset{a}{\to} p^+\pi^0$, have been considered in the literature since the sixties(Ginzburg & Zharkov 1965; Zharkov 1965; Ginzburg & Sirovatski 1965; Ritus 1985). The present available observational data on strongly magnetized astrophysical objects could provide an exciting scenario to test theoretical predictions involving such kind of inertially forbidden processes. The relevant observer-independent parameter characterizing these phenomena is the dimensionless quantity $\chi = a/m_p$, where $a$ and $m_p$ stand, respectively, for the proton proper acceleration and rest mass. (Unless otherwise stated, natural unities are adopted through this work.) For $\chi \ll 1$, one can employ a semiclassical approach where the proton is described by a classical current while the pion is considered as a fully quantized field. This corresponds to the so called no-recoil approximation, for which many analytical formulas are available. (For some recent works on the subject, see, for instance, (Herpay & Patkos 2008; Herpay et al. 2008) and (Fregolente, Matsas & Vanzella 2006; Fregolente & Saa 2008) for the electromagnetic and gravitational cases, respectively.) In particular, we expect a strong suppression of the pion emission for $\chi \to 0$ since such a process is well known to be inertially forbidden. For $\chi \gg 1$, a full quantized treatment is mandatory(Ritus 1985; Berezinsky, Dolgov & Kachelriess 1995). However, numerical evidences suggest that some appropriate limits of certain results obtained with the no-recoil approximation can still be considered as good estimates(Tokuhisa & Kajino 1999). For the case of the usual photon synchrotron radiation, for instance, quantum effects were carefully considered(Erber 1966) and the semiclassical approach was found to be accurate within a few percent in the limit $\chi \gg 1$. Notice that, in the cases where the proton acceleration is caused by strong magnetic fields, the parameter $\chi$ can be written in a similar way to the usual synchrotron radiation: $\chi = \gamma B/B_{\mathrm{cr}}$, where $\gamma$ is the Lorentz factor, $\gamma B$ is the magnetic field in the instantaneous reference frame of the proton, and $B_{\mathrm{cr}} = m_p^2/e \approx 1.5 \times 10^{20}$ G is a critical magnetic field strength, denoting the limit of the validity of the no-recoil approximation.

Thus, we have basically two regimes where the details of the pion emission by accelerated protons are known: the no-recoil regime $a \ll m_\pi$, where $m_\pi \approx 135$

---

Douglas Fregolente

Universidade Federal de Campina Grande, Centro de Formação de Professores, Cajazeiras, PB, Brazil

Alberto Saa

Departamento de Matemática Aplicada, UNICAMP, Campinas, SP, Brazil

[1] douglasfregolente@cfp.ufcg.edu.br

[2] asaa@ime.unicamp.br

MeV stands for the $\pi^0$ mass, and the ultra-accelerated one, $a \gg m_p \approx 938$ MeV. The intermediate acceleration range corresponding to $m_\pi < a < m_p$, which could encompass interesting astrophysical applications, is still open for debate. It is clear, however, that for such acceleration values, the pion emission channel will certainly be excited. In fact, the $\pi^0$ radiation process starts to become competitive with the usual photon synchrotron radiation only for $\chi \approx m_\pi/m_p$(Tokuhisa & Kajino 1999; Herpay et al. 2008).

The dominant decay mode for $\pi^0$ is the electromagnetic channel, giving origin to two photons(Nakamura et al. 2010). From the observational point of view, charged pions are more promising since they decay preferentially by the weak channel in muons and neutrinos(Nakamura et al. 2010). Processes like $p^+ \overset{a}{\to} n\pi^+$ have also been considered(Herpay & Patkos 2008; Fregolente, Matsas & Vanzella 2006), and they have essentially the same two regimes according to the value of the parameter $\chi$. We notice, however, that the semiclassical analysis of such processes in magnetic fields is rather tricky since the neutron will not be accelerated and, consequently, will follow a different trajectory from the proton, implying that the initial ($p^+$) and final ($n$) currents are qualitatively different, in sharp contrast to the gravitational case(Fregolente, Matsas & Vanzella 2006; Fregolente & Saa 2008). On the other hand, cases involving vector mesons like $p^+ \overset{a}{\to} p^+\rho$, with $m_\rho = 770$ MeV, as those ones considered in Tokuhisa & Kajino (1999), are analogous to the neutral $\pi^0$ case.

In the present work, we search the ATNF Pulsar Catalogue(Manchester et al. 2004) for strongly magnetized pulsars that could give rise to appreciable values for the parameter $\chi$. We employ rather conservative estimates based on the simplest polar cap model for pulsars. The highest value of $\chi$ we found corresponds to the 16 ms pulsar J0537-6910. Unfortunately, the obtained value $\chi \approx 10^{-6}$ is too small in order to give origin to observable signals. Nevertheless, our estimates reveal that a pulsar with period $P = 1\,\mathrm{ms}$ and surface polar magnetic field $B = 10^{12}$G would suffice to induce detectable pion radiation from accelerated protons. Our results can be applied also to the weak interaction channel emission, as in the process $p^+ \overset{a}{\to} ne^+\nu$. The threshold in this case correspond to the electron mass $m_e \approx 0.5$ MeV. The $\chi$ values for the known pulsars, however, are not enough to excite this emission channels either. For a $P = 1\,\mathrm{ms}$ pulsar, a magnetic field $B \approx 4 \times 10^{10}$G would be required to induce the weak interaction emissions.

## 2 Particle acceleration in the polar cap model

We use here the simplest polar cap model for pulsars. (For a comprehensive review on the subject, see Harding & Lai (2006).) We recall that, in the magnetosphere of strongly magnetized pulsars, one expects an electric field component parallel to the magnetic field, which could accelerate charged particles up to relativistic velocities along the magnetic field lines near the polar region of the star. Pulsars with polar caps for which $\mathbf{\Omega}\cdot\mathbf{B} < 0$, where $\mathbf{\Omega}$ and $\mathbf{B}$ stand, respectively, for the rotation and the magnetic field vectors, could effectively accelerate protons away from the polar region surface. In the so called inner acceleration region(Harding & Lai 2006), the accelerated particles are expected to radiate away their transverse energy so efficiently that they remain effectively constrained to move along the curved magnetic field lines with constant energy, corresponding to some Landau level.

The curvature radius $R_c$ of the polar magnetic field lines can be approximated by(Harding & Muslimov 1998)

$$R_c^2 \approx \frac{8G^2}{9\pi} rP, \tag{1}$$

where $r$ is the radius of the star, $P$ is the rotation period, and $G$ is a curved space correction term, which tends to increase $R_c$ by 25% - 30% when compared to the flat space formula. We assume the flat space value ($G = 1$) since we are going to estimate the maximum possible centripetal acceleration attained by protons. The curvature radius (1) does indeed correspond to the last open magnetic field line of the pulsar. For a relativistic particle following a circular trajectory with radius $R_c$, the proper centripetal acceleration is given by $a \approx \gamma^2/R_c$. Assuming that the kinetic energy $\epsilon_p$ of the accelerated proton moving along the last open magnetic field line was obtained from the parallel electric field, one has

$$\epsilon_p = \gamma m_p = \eta e \Delta\phi, \tag{2}$$

where $\eta$ accounts to the efficiency of the acceleration mechanism(Zhang & Harding 2000; Zhang et al. 2003) and

$$\Delta\phi = 2\pi^2 \frac{Br^3}{P^2} \tag{3}$$

is the Goldreich-Julian maximum potential drop near the star surface(Goldreich & Julian 1969), with $B$ being the pulsar surface polar magnetic field. The maximum efficiency is expect to be as high as 0.85 (Zhang & Harding 2000), but here we suppose total efficiency, $\eta = 1$. Finally, the maximum centripetal acceleration of such protons leaving the polar cap along the last open magnetic field line can be approximated by

$$\frac{a}{m_p} \approx 3\sqrt{2\pi^9} m_p \left(\frac{B}{B_{\rm cr}}\right)^2 \sqrt{\frac{r^{11}}{P^9}}. \tag{4}$$

For our purposes here, it is conveniente to introduce the usual dimensionless quantities $P_{-3}$ and $B_{12}$ defined, respectively, by $P = P_{-3} \times 10^{-3}$s and $B = B_{12} \times 10^{12}$G. Assuming a typical radius $r = 10^4$ m for the pulsar, one has finally

$$\chi = \frac{a}{m_p} \approx 0.35 B_{12}^2 P_{-3}^{-\frac{9}{2}}. \tag{5}$$

The total energy acquired by the proton is another important quantity in our analysis. From (2), we have

$$\gamma = 2\pi^2 m_p \frac{B}{B_{\rm cr}} \frac{r^3}{P^2}, \tag{6}$$

leading that

$$\gamma = 7 \times 10^9 B_{12} P_{-3}^{-2} \tag{7}$$

for stars of the typical size.

## 3 Results

We evaluate $\chi$ as given by (5) for all the 1700 pulsars in ATNF catalogue(Manchester et al. 2004) with known $P$ and $B$ values. Table 1 shows the ten pulsars with the highest values of $\chi$. As one can see, these values are too small to give rise to detectable signals, since the pion emission channel is effectively excited only for accelerations such that $\chi \approx m_\pi/m_p \approx 0.14$. Nevertheless, the proton can indeed reach considerable high energies, as the last column of Table 1 shows. For the J0537-6910 pulsar, for instance, the proton can reach an energy $\epsilon_p \approx 2.3 \times 10^5$ GeV. We notice that for the pulsar with the highest surface magnetic field (J1808-2024, $B \approx 2 \times 10^{15}$ G), one has only $\chi \approx 5.2 \times 10^{-12}$, since its period is considerable large ($P \approx 7.6$ s).

Figure 1 show the $(P_{-3}, B_{12})$ pairs for the considered pulsars. All the known pulsars are contained in the regions bounded by the curves (5) and (7) with, respectively, $\chi = 10^{-6}$ and $\gamma = 2 \times 10^7$ (upper) and $\chi = 10^{-14}$ and $\gamma = 2 \times 10^1$ (bottom). The small values obtained for $\chi$ are, of course, a consequence of the large proton mass $m_p$. Had we considered electrons instead of protons, we would have $B_{\rm cr} \approx 4.3 \times 10^{13}$G, and the parameter $\chi$ would be 10 orders of magnitude as larger. The regime $\chi \approx 1$ can be easily reached for electrons, explaining the rich physics of electrons in typical pulsar magnetospheres. Processes involving muons in circular orbits, like $\mu^- \to e^- \bar{\nu}_e \nu_\mu$ for instance, have also been investigated recently in the gravitational context (Fregolente & Saa 2008). The muon mass is $m_\mu = 107$ MeV, implying a muon acceleration parameter $\chi = a/m_\mu \approx 7.4 \times 10^{-4}$ for the pulsar J0537-6910. This is not enough to lead to any appreciable deviation of the muon half life as considered in Fregolente & Saa (2008).

Since we have $\chi \ll 1$, we can apply safely the formulas obtained in the no-recoil approximation for the process $p^+ \xrightarrow{a} p^+ \pi^0$. The total energy emitted by the pion synchrotron radiation, or in other words, the proton energy loss rate due to curvature pion radiation is given by(Tokuhisa & Kajino 1999)

$$\mathcal{R} = \frac{g^2}{\sqrt{3}} m_\pi m_p \chi \exp\left(-\frac{\sqrt{3}}{\chi}\frac{m_\pi}{m_p}\right) \tag{8}$$

in the limit $\chi \ll 1$, where $g^2 \approx 14$ is the strong coupling constant. The so called cooling time $t_\pi = \epsilon_p/\mathcal{R}$ is an important parameter corresponding to the time scale for which the process becomes relevant. For our case, we have

$$t_\pi = 6 \times 10^{-25} \gamma \chi^{-1} \exp\left(0.25\chi^{-1}\right) \text{ s}, \tag{9}$$

for stars of the typical size, with $\chi$ and $\gamma$ given, respectively, by (5) and (7). The exponential term in (9) gives rise to unreasonable high values of $t_\pi$ for the pulsars in the ATNP catalogue when compared, for instance, with the acceleration time $t_a$ (Herpay et al. 2008). From Figure 1, we can also infer the location of the curves with constant $t_\pi$ in the $(P_{-3}, B_{12})$ plane. Notice that the curves of constant $\chi$ and $\gamma$ are close in the log-log plot, since their inclination are, respectively 9/4 and 2. Regions of approximately constant $\chi$ and $\gamma$ would give rise, according to (9), to approximately constant $t_\pi$. So, the curve of constant $t_\pi$ passing through the point $(P_{-3}, B_{12})$ will be quite close to the lines of constant $\chi$ and $\gamma$ intercepting at the same point. For the the values of $\chi$ and $\gamma$ corresponding to the upper bound of the known pulsars, we would have unreasonable highs values for $t_\pi$. This is, of course, a consequence of the strong suppression of the pion radiation emitted by quasi-inertial protons. It is clear that no currently known pulsar could origin detectable signals of the curvature pion radiation emitted from accelerated protons in the magnetosphere.

**Acknowledgements** This work was supported by FAPESP and CNPq.



**Fig. 1** The values $(P_{-3}, B_{12})$ for the 1700 objects in the ATNF Pulsar Catalogue(Manchester et al. 2004) with known period $P = P_{-3} \times 10^{-3}$ s and surface magnetic field $B = B_{12} \times 10^{12}$ G. The uppermost red (solid) and green (dashed) lines correspond, respectively, to the curves of constant $\chi \approx m_\pi/m_p$ (5) and $\gamma \approx 4.5 \times 10^9$ (7). The other red (solid) and green (dashed) lines correspond, respectively, to the lines with $\chi \approx 10^{-7}, 10^{-9}, \ldots, 10^{-14}$ and with $\gamma \approx 10^7, 10^6, \ldots, 10^2$. All the pulsars in the ATNF catalogue are in a region of the plane $(P_{-3}, B_{12})$ from where no detectable signals are expected from curvature pion radiation emitted by accelerated protons in the magnetosphere.

| Pulsar | $P_{-3}$ | $B_{12}$ | $\chi_{-7}$ | $\gamma_7$ |
|---|---|---|---|---|
| J0537-6910 | 16.12 | 0.93 | 11.0 | 2.49 |
| B0531+21 | 33.08 | 3.78 | 7.26 | 2.42 |
| B0540-69 | 50.50 | 4.98 | 1.88 | 1.37 |
| J1813-1749 | 44.70 | 2.65 | 0.92 | 0.93 |
| J1400-6325 | 31.18 | 1.11 | 0.82 | 0.80 |
| J1747-2809 | 52.15 | 2.88 | 0.54 | 0.74 |
| J1833-1034 | 61.87 | 3.58 | 0.39 | 0.65 |
| J0205+6449 | 65.59 | 3.61 | 0.30 | 0.59 |
| J2229+6114 | 51.62 | 2.03 | 0.28 | 0.53 |
| J1617-5055 | 69.36 | 3.10 | 0.17 | 0.45 |

**Table 1** Top ten $\chi$ parameters, calculated according to (5), for the 1700 objects of the ATNF Pulsar Catalogue(Manchester et al. 2004) with known $P$ and $B$ values. The last column is the total proton energy given by (7). The pulsar period and magnetic field are, respectively, $P = P_{-3} \times 10^{-3}$s and $B = B_{12} \times 10^{12}$G. Analogously, we have $\chi = \chi_{-7} \times 10^{-7}$ and $\gamma = \gamma_7 \times 10^7$.

## References

Ginzburg, V. L., & Zharkov, G. F. 1965, Sov. Phys. JETP, 20, 1525.

Zharkov, G. F. 1965, Sov. J. Nucl. Phys., 1, 120.

Ginzburg, V. L., & Sirovatski, S. I., Ann. Rev. Astron. Astrophys., 3, 297.

Ritus, V. I. 1985, J. Sov. Laser Research, 6, 497.

Herpay, T., & Patkos, A. 2008, J. Phys. G, 35, 025201.

Herpay, T., Razzaque, S., Patkos, A., & Meszaros, P. 2008, JCAP, 0808, 025.

Fregolente, D., Matsas, G. E. A., & Vanzella, D. A. T. 2006, Phys. Rev. D, 74, 045032.

Fregolente, D., & Saa, A. 2008, Phys. Rev. D, 77, 103010.

Berezinsky, V., Dolgov, A., & Kachelriess, M. 1995, Phys. Lett. B, 351, 261.

Tokuhisa, A., & Kajino, T. 1999, Ap.J.Lett., 525, 117.

Erber, T. 1966, Rev. Mod. Phys., 38, 626.

Nakamura, K. *et al.* 2010 (Particle Data Group), J. Phys. G, 37, 075021 [http://pdg.lbl.gov]

Manchester, R. N., Hobbs, G. B., Teoh, A., & Hobbs, M. 2005, Astron. J., 129, 1993.

Harding, A. K., & Lai, D. 2006, Rept. Prog. Phys., 69, 2631.

Harding, A. K., & Muslimov, A. G. 1998, Astrophys. J., 508, 328.

Zhang, B., & Harding, A. K. 2000, Astrophys. J., 532, 1150.

Zhang, B., Dai, Z.G., Meszaros, P., Waxman, E., & Harding, A. K. 2003, Astrophys. J., 595, 346.

Goldreich, P., & Julian, W. H. 1969, Astrophys. J., 157, 869.

---

This manuscript was prepared with the AAS LaTeX macros v5.2.